

EUROPEAN PATENT APPLICATION

Application number: 89308232.1

Int. Cl.⁵: C12P 21/02 , C07K 1/00

Date of filing: 14.08.89

The title of the invention has been amended
(Guidelines for Examination in the EPO, A-III,
7.3).

Priority: 12.08.88 DK 4521/88

Date of publication of application:
21.03.90 Bulletin 90/12

Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

Applicant: CARLSBERG BIOTECHNOLOGY
LTD. A/S
Tagensvej 16
DK-2200 Copenhagen N(DK)


Inventor: Aasmul-Olsen, Stig

Skodsborgvej 410, st.tv.
 DK-2942 Skodsborg(DK)
 Inventor: Thorbek, Pia
 Sdr. Jagtvej 39
 DK-2970 Horsholm(DK)
 Inventor: Widmer, Fred
 c/o Carlsberg Biotechnology Ltd. A/S
 Tagensvej 16
 DK-2200 Copenhagen N(DK)
 Inventor: Hansen, Soren
 Tomsgaardsvej 102
 DK-2400 Copenhagen NV(DK)


Representative: Perry, Robert Edward et al
 GILL JENNINGS & EVERY 53-64 Chancery
 Lane
 London WC2A 1HN(GB)

Process for the enzymatic production of dipeptides.

Dipeptides or related compounds of the formula H-A-B-Y, wherein A is an L- or D- α -amino acid residue or ω -amino acid residue, B is an L- or D- α -aminocarboxylic acid residue, an L- or D-aminophosphonic acid residue or L- or D-aminosulfonic acid residue or the corresponding ω -amino acids or salts and hydrates thereof, and Y is OH, H, alkyl, aryl, aralkyl or a C-terminal blocking group, -or BY is an amino alcohol residue B¹-CHY¹-OH, wherein B¹ is a decarboxy derivative of the aminocarboxylic acid as defined with relation to B, and Y¹ is H, alkyl, aryl or aralkyl are produced by reacting a substrate component



wherein A is as defined above, R¹ is alkyl, aryl or aralkyl or an α -des-amino fragment of an amino acid, R² and R³ are hydrogen, alkyl, aryl or aralkyl with a nucleophile component selected from amino acids or their amides, esters, aldehydes or ketones of the formula



wherein B is an aminocarboxylic acid residue as defined above, R² and R³ have the above meaning, or

when R^2 is hydrogen, R^3 is also hydroxy or amino, R^4 and R^8 are alkyl, aryl or aralkyl, and may also be H, aminophosphonic acids or aminosulfonic acids

$NH_2C_xH_zPO_3R^5R^6$ or $NH_2C_xH_zSO_3R^7$

wherein R^5 , R^6 and R^7 are hydrogen, alkyl, aryl or aralkyl, x is 1-6 and z is 2-12, and

amino alcohols $H-B'-CHY'-OH$ wherein B' and Y' are as above

in the presence of an amidase or esterase enzyme different from serine or thiol carboxypeptidases, preferably a serine or thiolendoprotease, lipase, esterase or glycosidase in solution or dispersion, and then, if desired, cleaving a present side-chain protecting group or protective group Y and/or, if desired, converting the resulting dipeptide derivative to a salt or hydrate.

The process allows for production of LL-, LD-, DL- and DD-dipeptides without risk of racemization in a simple and economic manner.

A process for enzymatic production of dipeptides or structurally related compounds

The present invention concerns a process for enzymatic production of dipeptides and derivatives of dipeptides and having the general formula

H-A-B-Y

wherein A represents an optionally side-chain protected L- or D- α -amino acid residue or ω -amino acid residue and B represents an optionally side-chain protected L- or D- α -aminocarboxylic acid residue which may be the same as or different from A, and L- or D-aminophosphonic acid residue or L- or D-aminosulfonic acid residue or the corresponding ω -amino acids or salts and hydrates thereof, and Y is OH, H, alkyl, aryl, aralkyl or a C-terminal blocking group, or BY represents an amino alcohol residue B'-CHY'OH

wherein B' is a decarboxy derivative of the aminocarboxylic acids as defined with relation to B, and Y' is H, alkyl, aryl or aralkyl.

In recent years, there has been an increasing interest in dipeptides and dipeptide derivatives optionally containing an amino acid residue of D-configuration, with a view to their potential pharmacological effects, such as e.g. antibiotics. Likewise, there has been a great interest in dipeptides within fields such as artificial nutrition -human as well as veterinary -sweeteners and within agrochemistry, such as e.g. herbicides.

Such dipeptides H-A-B-Y can be produced by means of known chemical coupling reactions, but all these methods share the feature that, generally, it is necessary to protect the amino acids involved - A and B - on the amino group and the carboxylic acid group, respectively, and frequently also on the side chains if these carry functional groups. Further, there is an inherent risk of side reactions during the chemical coupling step because of the reagents and conditions employed, a major side reaction being racemization, particularly of the A-component. By replacing the chemical coupling step with an enzymatic coupling step proceeding under mild conditions, such side reactions and racemization can be avoided, yielding a stereochemically pure product.

The presence of amino- and carboxyl protective groups is mandatory in chemical coupling and has generally been regarded as mandatory also in prior art enzymatic coupling using endoproteases.

This adds several undesired features to these processes seriously afflicting their process economy on an industrial scale, particularly apparent in dipeptide synthesis.

The disadvantages are concerned with the introduction of these groups, as well as their removal and presence during process operation, increasing overall process cost and time and affecting overall yield.

Typical examples of amino protective groups commonly used are those of the carbobenzoxy (Z-) and tert-butoxycarbonyl (Boc-) type, which are of a molecular weight comparable to those of the amino acid residues. Firstly, the protective groups will have to be introduced in the starting materials by means of appropriate costly agents in a separate reaction step followed by an isolation step. While present, these hydrophobic groups often have a drastical effect upon the solubility of the intermediates and reaction products, and may afflict both the nature and the amount of solvents required in their processing as well as ease of purification and of deprotection. The deprotection will also take place in a separate step with a following purification step.

For this purpose a series of reactions are available, but with the exception of catalytical hydrogenation, posing industrial problems of its own, these methods are occurring under violent, often strongly acidic or basic conditions, frequently giving rise to a series of side reactions, resulting in an impure product or demanding laborious purification.

The last steps in this relatively long series of synthesis steps may thus be a rather comprehensive deprotection to obtain the desired peptides, and, owing to the almost inevitable secondary reactions, rather laborious purification procedures are frequently required to provide a product with the desired high purity.

A dipeptide which has attracted great attention in recent years is L-Asp-L-Phe-methylester, also known as aspartame, and derivatives thereof which has found extended use as sweeteners. The chemical synthesis of aspartame is entailed with the above-mentioned drawbacks.

Attempts to avoid amino terminal protection in the production of aspartame and its derivatives have led to microbial fermentation approaches, like the fermentation processes described in EP-A1-074095, EP-A2-102529 and EP-A2-154472. This technique is fundamentally different from synthetic approaches and relies on specific organisms for aspartame, and is thus not generally applicable in connection with other dipeptides. In addition, the yields are low and recovery from the fermentation broth laborious.

The above-mentioned shortcomings in the known processes for the production of aspartame are confirmed in EP-A2-269390. In this application a method for producing L-Asp-L-Phe-alkylesters is claimed which comprises reacting in a solvent medium L-aspartic acid alpha ester or L-aspartic acid alpha amide

with L-phenylalanine alkyl ester in the presence of an enzyme, microorganism containing the enzyme, enzyme containing fraction of a microorganism, or enzyme immobilized on solid support, said enzyme being capable of forming L-aspartyl-L-phenylalanine alkyl ester by condensation of the L-aspartic acid alpha ester or L-aspartic acid alpha amides and L-phenylalanine alkyl ester.

5 It is seen that an enzyme must be used which has specific esterase or amidase activity against Asp, but not against Phe. The only enzyme mentioned is the extracellular protease with esterolytic activity of *Staphylococcus aureus* V8.

While therefore the applicability of N- α -unprotected Asp-esters or amides is suggested - along with N-protected or β -esters or amides - the disclosure of one specific enzyme for the use in the production of a
10 specific peptide supported only by one example where rather peculiar reaction conditions are used, viz. 5 times excess of substrate components, this reference does not by far provide a general teaching of the applicability of N-unprotected substrate components in dipeptide synthesis catalyzed by amidase or esterase enzymes.

Thus, it is an obvious advantage in terms of overall process economy to be able to avoid protective
15 groups, also on the amino and carboxy terminus. In some cases, it may be of interest to be able to produce a dipeptide having side-chain protection, but no terminal protection, and it will be shown that it is possible in the process according to this invention, starting from side-chain protected, but amino unprotected starting materials. In this case, the same advantages of mild reaction conditions and overall process economy may be obtained. If desired, the side-chain protective group may be removed by chemical or enzymatic means,
20 generally under milder conditions than amino protective groups.

The enzyme catalyzed coupling reactions enabling the use of optionally side-chain unprotected amino acid derivatives and an optionally C-terminal unprotected B-component (nucleophile) are known. See e.g. the DK Patent Specification No. 155613 as well as the corresponding EP Patent Specification No. 17 485 (EP-B1-17485), incorporated herein by reference.

25 Briefly, EP-B1-17485 describes a process for producing peptides by reacting a substrate component selected from i.a. amino acid esters and amino acid amides with an amine component (nucleophile) selected from i.a. L-amino acid amides, L-amino acids or L-amino acid esters in the presence of an L-specific serine or thiol carboxypeptidase enzyme.

If a dipeptide is to be produced by the process of EP-B1-17485, the substrate component is obligatory
30 an N-terminal protected amino acid derivative, and the constituent amino acid is obligatory an L-amino acid.

As described in EP-A1-278787, (incorporated by reference) and other applications claiming priority from the DK appln. 725/87 filed on February 13, 1987 it was surprisingly found that the serine and thiol carboxypeptidases used in EP-B1-17485 are capable of utilizing N-unprotected amino acid esters or amides as a substrate component in controlled reactions for synthesis of dipeptides and dipeptide derivatives, and
35 that it was possible to suppress a possible oligomerization of the substrate.

Admittedly, it has long been known that some endoproteases can catalyze oligomerization of certain N-unprotected amino acid esters with L-configuration, (Fruton, J.S. *Advances in Enzymology*, 53: 239-306, 1982) but it has never been attempted to use this for production of dipeptides which are not simple dimers. Generally, the results of such oligomerizations have been a mixture of a series of oligomers, sometimes
40 long. The degree of oligomerisation and complexity of the mixture depend on the solubility of the products formed. Also it should be mentioned that the mere fact that a given enzyme can hydrolyze a particular N- α -unprotected amino acid derivative does not automatically lead to a similar ability to catalyze coupling reactions involving the same derivative. See the references cited by Andersen, A.J. in "Peptides, Structure and Function". *Proc. of the ninth ann. peptide symposium*, Eds. Deber, C.M. et al., Pierce (1985), p. 355.

45 For this reason the use of serine and thiol endoproteases for peptide synthesis has been limited to the use of amino and carboxy terminal protected starting materials, as exemplified by US-A-4.086.136, where e.g. papain, strombromelein, ficin, chymopapain and chymotrypsin are mentioned.

Furthermore, for these enzymes free amino acids have so far generally been regarded as unsuitable as amino components as noted by Y.W. Mitin et al., *Int. J. Peptide Protein Res.*, Vol. 23 (1984), p. 528-534.

50 The above-mentioned amino and carboxy terminal protected starting materials are also mandatory if aspartate endoproteases e.g. pepsin are used as exemplified by US-A-3.972.773, and if metallo endoproteases are used, as exemplified by the synthesis of Z-AspPheOMe.PheOMe-salt in EP-A1-009585. In these condensation type reactions it is further mandatory that the substrate component is on the free α -carboxylic acid form, as exemplified in EP-A2-220923.

55 The use of amino and carboxy terminal protected starting materials has also been considered mandatory for nonproteases, e.g. esterases with amidase activity, Blair West et al., *Tetrahedron letters*, Vol. 28. No. 15, p. 1629-32, (1987), who used porcine pancreatic lipase, *Candida cylindracea* lipase and pig liver esterase in organic media, optionally containing an aqueous buffer.

Cellulases in turn, have so far only been used synthetically for polymerisation condensation type reactions of completely free β -amino acids as described by Kitazume, T. et al., J. Flourine Chem. 36 - (1987), p. 225-236.

The synthesis of the diastereomeric dipeptides of DL, LD and DD-configuration as well as peptides containing β -amino acid residues from amino-unprotected substrate components has so far not been possible with carboxypeptidases except as described with relation to EP-A1-278787 nor in general with any proteolytical enzymes (Class EC 3.4). Some efforts have been made with a different class of enzymes, aminoacyl-t-RNA-synthetase (Class LC 6.1) as exemplified by EP-A1-086053. In this case a specific enzyme must be used for each type of amino acid residue, and furthermore, expensive Co-factors like ATP are required. At the same time, yields are very poor, so even though some product was isolated and identified, typically a ten fold excess of Co-factor and a hundred fold excess of nucleophile and up to a thousand fold excess of enzyme by weight was required.


A recent study by Gaertner et al., Proteins: Structure, Function and Genetics 3: 130-137 (1988) indirectly confirms the prejudice against using N- α -unprotected amino acids as substrate components in enzymatic peptide synthesis. Since a major problem earlier encountered is secondary hydrolysis of the newly synthesized product, an attempt was made to decrease water activity in the reaction medium by using organic solvents. Gaertner et al. reports dipeptide synthesis using chymotrypsin which had been chemically modified to enhance its solubility in organic media. Using a number of N- α -protected amino acid esters as substrates and amino acid amides as nucleophiles N- α -protected dipeptide amides were obtained in good yields in a benzene medium. Thus, α -Bz-Lys-Phe-NH₂ was obtained in a yield of more than 98% by reacting α -Bz-Lys-OMe with Phe-NH₂. However, when using ϵ -Z-LysOEt, i.e. a side-chain protected but N- α -unprotected lysine ester, no ϵ -Z-Lys-Phe-NH₂ was formed. Gaertner et al do not comment on this result, but it indicates that they apparently believed that amino group protection of the substrate component was compulsory.

The present invention which represents a further development of the invention according to EP-A1-278787 resides on the surprising finding that the capability of utilizing N- α -unprotected amino acid esters and amides of the substrate component in controlled reactions for synthesis of dipeptides and dipeptide derivatives is not limited to serine or thiol carboxypeptidases, but also possible with amidase or esterase enzymes generally, in particular serine endoproteases, thiol endoproteases, lipases and esterases.

Again, it was surprisingly found that also N- α -unprotected amino acid derivatives of D-configuration can be used as substrates in these reactions, so that, in addition to LL-dipeptides, it is also possible to synthesize DL-dipeptides. The reaction rate for D-substrates, however, is generally somewhat lower than for L-substrates under uniform conditions, but the difference in rate is much smaller than for the corresponding N-protected amino acid esters, the D-substrate being reacted at a rate which is much smaller than the rate for the L-substrate; Purdie et al., Biochem. Biophys. Acta, 268 (1972), 523. There are, however great individual differences between the enzymes in this respect. Synthesis yields are often just as high with the unprotected D-substrates in relation to the unprotected L-substrates.

It is known that certain endoproteases are capable of utilizing a nucleophile component having D-configuration in reactions with N-protected substrate components. Thus, depending on the actual sequence it is possible to synthesize dipeptides having LL-, DL-, LD- or DD-configuration by the process according to the invention.

The process of the invention is thus characterized by reacting a substrate component, which is an amino acid derivative having the formula




wherein A is as defined above, R¹ represents alkyl, aryl or aralkyl optionally substituted with inert substituents or an α -des-amino fragment of an amino acid, and R² and R³ are the same or different and each represents hydrogen, alkyl, aryl or aralkyl optionally substituted with inert substituents, with a nucleophile component selected from

(a) amino acids having the formula



wherein B is an aminocarboxylic acid residue as defined above,

(b) amino acid amides having the formula




wherein B is an aminocarboxylic acid residue as defined above, and R² and R³ have the above meaning, except that when R² represents hydrogen, R³ may also represent hydroxy or amino.

(c) amino acid esters having the formula




wherein B is an aminocarboxylic acid residue as defined above, and R⁴ represents alkyl, aryl or aralkyl.

(d) optionally acid group protected straight chain or branched aminophosphonic acids or aminosulfonic acids having the formula




wherein R⁵, R⁶ and R⁷ independently represent hydrogen, alkyl, aryl or aralkyl, x is 1-6 and z is 2-12.

(e) amino acid aldehydes or ketones or derivatives thereof having the formula



wherein B' is as defined above, Y² is O or a functional derivative thereof, preferably a ketal, and R⁸ is H, alkyl, aryl or aralkyl, and

(f) amino alcohols having the formula



wherein B' and Y' have the above meaning in the presence of an amidase or esterase enzyme different from serine or thiol carboxypeptidases in solution or dispersion, and then, if desired, cleaving a present side-chain protecting group or protective group Y and/or, if desired, converting the resulting dipeptide derivative to a salt or hydrate.

Examples of useful amino acids include aliphatic amino acids, such as monoaminomonocarboxylic acids, e.g. glycine (Gly), alanine (Ala), valine (Val), norvaline (Nval), leucine (leu), isoleucine (iso-Leu) and norleucine (Nleu), hydroxy amino acids, such as serine (Ser), threonine (Thr) and homoserine (homo-Ser), sulfur-containing amino acids, such as methionine (Met) or cystine (CysS) and cysteine (CysH), monoaminodicarboxylic acids, such as aspartic acid (Asp), glutamic acid (Glu) and amides thereof, such as asparagine (Asn) and glutamine (Gln), diaminomonocarboxylic acids, such as ornithine (Orn) and lysine (Lys), arginine (Arg), aromatic amino acids, such as phenylalanine (Phe) and tyrosine (Tyr), as well as heterocyclic amino acids, such as histidine (His), proline (Pro) and tryptophan (Trp). As examples of useful amino compounds of a more unusual structure may be mentioned penicillamine (Pen), aminophosphonic acids, such as alanine-phosphonic acid (AlaP), aminosulfonic acids, such as taurine (Tau), ω -amino acids, such as β -alanine (BAla), iso amino acids, such as α -methylalanin (Aib), amino acids substituted with inert substituents, e.g. halogen or nitro, or the aldehyde, ketones, ketals or alcohols derived from e.g. alanine. As mentioned, they may be included in D-form in the substrate component and they may also be present in D-form in the nucleophile component.

It will be understood that the definitions given for the group Y in the peptide derivative of the formula H-A-B-Y are reflected in the various nucleophile components, stated in claim 1. Thus, (b) and (c) both contain C-terminal blocking groups, while Y = H corresponds to an amino acid aldehyde and Y = alkyl, aryl or aralkyl corresponds to an amino ketone. If desired a functional derivative of an amino ketone may be used as nucleophile, e.g. a ketal, oxime or sulfite.

As in the process according to EP-A1-278787, the advantages of the process of the present invention over the mentioned known methods are minimum or no side chain protection, no N-protection of the substrate component which may have both D- and L-configuration, no risk of racemization, few synthesis steps and an expected relatively pure end product, which in combination provides an extremely simple and economic method of production.

Preferred substrate components are esters in which R¹ is a straight or branched alkyl having 1 to 8 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, amyl, hexyl, heptyl and octyl, or the aralkyl group benzyl. Particularly expedient nucleophile components are amino acid amides, in which R² is H, and R³ is H or C₁-C₆ alkyl, or amino acid esters in which R⁴ is a straight or branched alkyl having 1 to 6 carbon atoms such as the above-mentioned ones. As mentioned, R¹ may be alkyl, aryl or aralkyl optionally substituted with inert substituents, e.g. hydroxy or nitro.

The invention also comprises the processes involving intermediate formation of a peptide containing the group



5 following which this group may be cleaved to form a carboxylic acid group. This cleavage may be catalyzed by another enzyme or the same enzyme as was used to form the peptide, albeit under different reaction conditions. Also C-terminal modifications of the group Y may be made.

10 Enzymes may also be used to cleave side-chain protective groups, applicable enzymes being proteolytic enzymes, lipases and esterases, according to the nature of the protective group, see "The peptides, Analysis, Synthesis, Biology" Vol 9, Special Methods in peptide Synthesis Part C. J.A. Glass, Enzymatic manipulation of Protecting Groups in Peptide Synthesis, Academic Press 1987.

15 As examples of enzymes possessing esterase and/or amidase activity, and therefore expected to be active as catalysts in the process according to this invention may be mentioned those listed in the table below.

For a closer description of these enzymes, reference is made to i.a. Perlmann, G. E. et al., in Colowick, S. P. & Kaplan, N. O. (eds.) Methods of Enzym. 8 (1966) 19 (1970), 28 (1972), 35 (1975) and 45 (1976), Fruton, S., Adv. Enzymol. 53, p. 239, Wiley (1982), Abassi, A. et al. Biol. Chem. Hoppe-Seyler, 367, p. 441-45 (1986), Dixon, M. and Webb E. C. "Enzymes", 3rd ed. Longman Group (1979), Torrey, S. (ed.) in "Enzyme Technology, Recent Advances", Biotech, Rev. 2, Noyes Data Corporation (1983), and finally Laane, C., Tramper, S., Lilly, M. D. (eds.) "Biocatalysis in Organic Media", Elsevier (1987), Asano, Y. et al., Angew. Chem. 101, (1989), p. 511-512, Kitazume, T. et al., J. Fluorine Chem. 36 (1987), p. 225-236. all being incorporated by reference.

TABEL II. Proteases different from carboxypeptidases:

| | Enzyme | (Abb) | Normal Source |
|--|--|-------|---|
| A) <u>Thiolendo-</u> <u>proteases:</u> | Papain | (P) | Papaya |
| | Chymopapain | (CP) | Papaya |
| | Bromelain | (B) | Pineapple |
| | Ficin | (F) | Fig (Ficus) |
| | Clostripain | (CL) | Clostridium histolyticum |
| B) <u>Serineendo-</u> <u>proteases:</u> | Trypsin | (T) | Pancreas |
| | Chymotrypsin | (CT) | Pancreas |
| | Elastase | (E) | Pancreas |
| | Subtilisin | (S) | Bacillus licheniformis or subtilis |
| | Thermitase | (TV) | Thermoactino- myces vulgaris |
| | Proteinase K | (K) | Tritirachium album |
| | Valyl-proteinase | (VP) | Candida tropicalis |
| | Post Prolin Specific Endopeptidase (PPSE) | | Flavo- bacterium meningo- septicum |

5 Achromobacter lyti-
cus Protease I (AL-I) Achromobacter
lyticus
10 Endoproteinase ArgC (AC) Submaxil-
laris glands
Endoproteinase LysC (LC) Lysobacter
enzymogenes
15 Thrombin (TB) Blood plasma

C) Exopeptidases:

20 Aminopeptidase (EC3.4.11) Animal,
particularly from vegetable,
Achromobacter sp. or microbial

25 II. Other hydrolases acting on ester bonds

Carboxylesterase (EC.3.1.1.1) Liver
30 Arylhydrolase (EC.3.1.1.2) Plasma
Triacylglycerollipase (EC.3.1.1.3) Animal,
particularly from vegetable,
35 Porcine Pancreas or or micro-
Candida cylindracea or bial
Lipolase* (NOVO); (Aspergillus sp.)
40 Acetic ester
acetylerase (EC.3.1.1.6) Animal
tissue
45 Arylglycerol lipase (EC.3.1.1.23) Animal or
vegetable

III Glycosidases

50 Cellulase (EC.3.2.1.4) Animal,
particularly from vegetable,
Trichoderma viride or or micro-
55 Aspergillus niger bial

The presently preferred enzymes are trypsin, chymotrypsin, subtilisin, elastase, papain, chymopapain, clostripain, porcine pancreatic lipase and *Candida cylindracea* lipase.

The enzyme used may also be chemically modified or be a biosynthetic mutant of a natural form.

5 As illustrated more fully below, the process of the invention is rather simple.

The reaction may be performed in an aqueous reaction medium, if desired containing up to 90%, preferably up to 60% of a polar organic solvent which is miscible with water, and compatible with the enzyme under the conditions specified. Preferred solvents are lower alcohols, dimethyl formamide, dimethyl sulfoxide, dimethoxy ethane and ethylene glycol.

10 It is important to maintain a rather constant pH value in the reaction mixture. This pH value is between 3 and 11, preferably between 5 and 10.5, more preferably between 6 and 10 and most preferably between 7 and 9.5 and also depends upon the concrete starting materials, the peptide formed and the enzyme.

The reaction may alternatively be performed in a nonaqueous, nonpolar medium being composed of water immiscible organic solvents e.g. benzene, toluene, hexanes, heptanes, octanes, dialkyl ethers, ethyl acetate, ethyl propionate and methylene chloride, etc. which are compatible with enzymes having esterase
15 and/or amidase activity, in particular lipases.

These essentially apolar reaction media may contain up to 10% water in dissolved form in order to facilitate optimal enzyme performance or reaction stability.

The reaction temperature is preferably room temperature and above, 20 to 50 °C, but temperatures in the range 0 to 80 °C may be used, if advantageous under the conditions otherwise given. At high solvent conditions subzero or temperatures higher than 80 °C may be used.

The concentration of the two reaction components may vary within wide limits, but the nucleophile component is frequently in excess, and to avoid oligomerization of the substrate component, said component is often added in minor portions at intervals during the entire reaction sequence. If a good
25 nucleophile for the enzyme is used in high excess, surprisingly no oligomerization is observed, and high substrate concentrations may be used without side reactions. In case a homo-dipeptide in which A = B is wanted, oligomerization from a single nucleophile can be avoided by using different carboxy protective groups on the substrate and nucleophile component.

Thus, the starting concentration of the substrate component may typically be 0.005 to 2 molar and for
30 the nucleophile component in cases, where it is added separately, 0.005 to 3 molar. In most situations it is possible to recover excess of the nucleophile component and the hydrolysis product from the substrate component for optional reesterification and reuse. Recycling of the components is particularly easy because of their simple structure and the absence of side reactions and deprotective losses.

The enzyme concentration may likewise vary, but is frequently somewhat higher (5-50 µm) (or even
35 higher, up to ~ 500 µm) than the concentrations appropriate in the use of N-protected amino acid ester substrates, but the amount required for synthetic purposes may be reduced more than tenfold by using a stable immobilized enzyme preparation, thereby enabling the enzyme to be used in a continuous process.

The reaction medium may also contain salts, such as NaCl and CaCl₂, which influences the binding of the enzyme to the substrate and may stabilize the enzyme, as well as a complex binding agent for present
40 metal ions, such as EDTA, and mercaptostabilizing agents such as DTT, BME or Cysteine may be used with for instance thiolendoproteases.

This surprising hydrolysis sequence is reflected in the following examples, which illustrate the production of various dipeptides in the process of the invention using various enzymes.

45

General method for examples 1-13 and 20 et seq.

The reactions, performed on an analytical scale with a reaction volume of 1 ml, were carried out in a pH-stat, and the selected pH value was kept constant by automatic addition of 1 N NaOH except for the
50 examples with high content of organic solvents or immobilized enzymes, where the conditions are indicated in the individual examples. Reaction temperature was room temperature, unless otherwise stated. The table also includes reaction concentrations, content of organic solvent, product and yield. Reaction times are typically between 0.5 and 5 hours and the enzyme concentrations are typically 5-20 µm, unless otherwise stated.

55 Product identification and determination of product yield were performed by means of reverse phase HPLC (Waters 6000 A pumps, 660 gradient blender, UK 6 injector) on a C₁₈NOVA PAK column (Waters, RCM) using suitable gradients of elution systems containing 50 mM triethylammonium phosphate, pH 3.0 from 0% to 80% acetonitrile with a flow of 2 ml/min. Elution was monitored by means of a UV detector

(Waters 480) at 230 nm, 254 nm, 278 nm or 290 nm.

The products were identified by amino acid analysis of fractions from the HPLC analysis, which corresponded to the assumed product peak and/or by HPLC comparison with a chemically synthesized reference product. These were produced according to known principles, usually via reaction between BOC-A-OSu - the tertiary butyloxy carbonyl - succinic imide ester derivative of the substrate amino acid - and the used nucleophile component followed by deblocking of the N-terminal amino acid residue. In all cases, it was possible to separate LL-and DD-dipeptides from the diastomeric DL-and LD-dipeptide products.

For the products which can only be detected at 230 nm, the product yields were determined by means of the absorption/concentration curve of the chemically synthesized reference compound. For the other products, the yields were determined on the basis of the ratio between the integrated areas below the peaks in the elution chromatogram, corresponding to the product respectively the reactant which absorbs at the wavelength concerned.

The reaction conditions in the preparative examples 14-17 are described in the individual examples. The reactions were followed on analytical HPLC as described. The enzyme concentrations are generally lower and the reaction times longer than in the corresponding analytical examples, but no attempt to optimize the reaction conditions has been made.

| Example 1 | | | |
|---|----------|-----------------------|-------|
| Trypsin ^{a)} catalyzed synthesis of L,L-dipeptide amides with L-Arginine ethyl ester (50 mM) as substrate and L-amino acid amides as nucleophiles at various concentrations in water at pH 8.5 | | | |
| Nucleophile | (conc.) | Product | Yield |
| Leucine amide | (0.2 M) | ArgLeuNH ₂ | 20% |
| Leucine amide | (0.7 M) | ArgLeuNH ₂ | 32% |
| Methionine amide | (0.25 M) | ArgMetNH ₂ | 31% |
| Methionine amide | (0.5 M) | ArgMetNH ₂ | 52% |
| Methionine amide | (1.0 M) | ArgMetNH ₂ | 90% |
| Serine amide | (0.5 M) | ArgSerNH ₂ | 45% |
| Tyrosine amide | (0.5 M) | ArgTyrNH ₂ | 46% |

^{a)} 10 uM

| Example 2 | | | | |
|--|--------------------|---------|-----------------------|-------|
| Trypsin ^{a)} catalyzed synthesis of diastereomeric L,D- and D,L-dipeptide amides with L- or D-Arginine ethyl ester (50 mM) as substrate and L- or D-Leucine and Methionine mides as nucleophiles in water at pH 8.5 | | | | |
| Substrate | Nucleophile | (conc.) | Product | Yield |
| D-argOEt | L-Leucine amide | (0.2 M) | argLeuNH ₂ | 20% |
| D-argOEt | L-Leucine amide | (1.0 M) | argLeuNH ₂ | 30% |
| D-argOEt | L-Methionine amide | (0.5 M) | argMetNH ₂ | 68% |
| L-ArgOEt | D-methionine amide | (1.0 M) | ArgmetNH ₂ | 5% |

^{a)} 10 uM

Example 3

Trypsin ^{a)} catalyzed synthesis of L,L-dipeptide amides using L-amino acid amides (0.5 M) as nucleophiles and L-Lysine or Histidine ethyl ester (50 mM) as substrate in water at pH 8.5

| Substrate | Nucleophile | Product | Yield |
|-----------|-------------------|-----------------------|-------|
| LysOEt | Methionine amide | LysMetNH ₂ | 81% |
| LysOEt | Tryptophane amide | LysTrpNH ₂ | 32% |
| LysOEt | Alanine amide | LysAlaNH ₂ | 5% |
| HisOEt | Methionine amide | HisMetNH ₂ | 62% |

^{a)} 10 μ M

Example 4

Alpha-Chymotrypsin ^{a)} catalyzed synthesis of L,L-dipeptide amides using L-Tyrosine ethyl ester (50 mM) as substrate and L-amino acid amides as nucleophiles in water

| Nucleophile | (conc.) | pH | Product | Yield |
|----------------|---------|-----|-----------------------|-------|
| Leucine amide | (0.2 M) | 9.0 | TyrLeuNH ₂ | 68% |
| Arginine amide | (0.4 M) | 8.5 | TyrArgNH ₂ | 90% |
| Serine amide | (0.4 M) | 8.5 | TyrSerNH ₂ | 75% |

^{a)} 5 μ M

Example 5

Alpha-Chymotrypsin ^{a)} catalyzed synthesis of L,D-dipeptide amides using L-Tyrosine ethyl ester (5 or 50 mM) as substrate and D-amino acid amides as nucleophiles in water

| Nucleophile | (conc.) | pH | Product | Yield |
|--------------------|---------|-----|-----------------------|-------------------|
| D-Leucine amide | (0.2 M) | 9.0 | TyrLeuNH ₂ | 17% ^{b)} |
| D-isoleucine amide | (0.3 M) | 9.0 | TyrIleNH ₂ | 23% ^{b)} |
| D-serine amide | (0.4 M) | 8.5 | TyrSerNH ₂ | 35% |

^{a)} 5 μ M

^{b)} 5 mM substrate

Example 6

Alpha-Chymotrypsin catalyzed synthesis of D,L-dipeptide amides using D-tyrosine ethyl ester (50 mM) as substrate and L-Leucine amide as nucleophiles in water at pH 9.0

| Nucleophile | (conc.) | Product | Yield ^{c)} |
|-----------------|---------|---------------------------|---------------------|
| L-Leucine amide | (0.2 M) | D,L-tyrLeuNH ₂ | 40% ^{a)} |
| L-Leucine amide | (0.3 M) | D,L-tyrLeuNH ₂ | 68% ^{b)} |

^{a)} 50 uM enzyme

^{b)} 100 uM enzyme

^{c)} Prolonged reaction time - days

Example 7

Alpha-Chymotrypsin ^{a)} catalyzed synthesis of L,L- and L,D-dipeptide amides and esters using different L-amino acid esters (50 mM) as substrates and L- or D-amino acid esters (0.8 M) or amides as nucleophiles at pH 8.5

| Substrate | Nucleophile | Solvent | Product | Yield ^{d)} |
|-----------|------------------------------------|---------|-----------------------|---------------------|
| L-PheOEt | L-ArgNH ₂ ^{b)} | Water | PheArgNH ₂ | 82% |
| L-TyrOEt | L-SerOEt | Water | TyrSerOEt | 48% ^{c)} |
| L-TyrOBzl | L-SerOEt | 30% DMF | TyrSerOEt | 40% ^{c)} |
| L-TyrOBzl | L-SerOMe | 30% DMF | TyrSerOMe | 39% ^{c)} |
| L-TyrOBzl | D-serOMe | 30% DMF | TyrserOMe | 21% ^{c)} |

^{a)} 5 uM

^{b)} 0.4 M nucleophile

^{c)} at 90% conversion

^{d)} hydrolysis/diketopiperazine formed due to chemical instability of product was observed under these conditions in amounts of 15-35% and is not included in the yields reported.

Example 8

Subtilisin A ^{a)} catalyzed synthesis of sidechain protected L-Aspartyl-D-alanyl amide using L-Aspartyl diesters (0.1 M) as substrates and D-alanine amide as nucleophile at pH 8.5

| Substrate | Nucleophile (conc.) | Solvent | Product | Yield |
|--|---------------------|----------|-----------------------------|-------|
| L-Asp(OEt) ₂ | (1.0 M) | Water | Asp(OEt)alaNH ₂ | 9% |
| L-Asp(OEt) ₂ | (2.0 M) | Water | Asp(OEt)alaNH ₂ | 24% |
| L-Asp(OBzl) ₂ ^{b)} | (0.5 M) | 30% DMSO | Asp(OBzl)alaNH ₂ | 8% |
| L-Asp(OBzl) ₂ ^{b)} | (1.0 M) | 30% DMSO | Asp(OBzl)alaNH ₂ | 20% |

^{a)} 5 uM

^{b)} 50 mM substrate, 20 uM enzyme

Example 9

Elastase ^{a)} catalyzed synthesis of L,L-amides using amino acid benzyl ester (50 mM) as substrate and amino acid amide (0.5 M) as nucleophile in water at pH 8.5

| Substrate | Nucleophile | Product | Yield |
|-----------|------------------|-----------------------|-------|
| L-ValOBzl | L-Arginine amide | ValArgNH ₂ | 59% |

^{a)} 40 uM

Example 10

Synthesis of L,L-dipeptide amides catalyzed by papaya thiolendoproteases^{a)} using amino acid esters (50 mM) as substrates and L-amino acid amides (0.8 M) as nucleophiles in water at pH 8.5

| Enzyme | Substrate | Nucleophile | Product | Yield |
|-------------|-----------|--------------------|-----------------------|-------|
| Papain | LysOEt | AlaNH ₂ | LysAlaNH ₂ | 60% |
| Papain | LysOMe | AlaNH ₂ | LysAlaNH ₂ | 55% |
| Chymopapain | LysOEt | AlaNH ₂ | LysAlaNH ₂ | 23% |
| Chymopapain | LysOMe | AlaNH ₂ | LysAlaNH ₂ | 34% |

^{a)} 100 uM, 2mM EDTA, 10 mM Cysteine

Example 11

Clostripain ^{a)} catalyzed synthesis of L,L-dipeptide amides and esters with L-Arginine ethyl ester as substrate (50 mM) and L-amino acid amides as nucleophiles at pH 8.5

| Nucleophile | (conc.) | Solvent | Product | Yield |
|-----------------------|---------|----------|-----------------------|-------|
| L-Methionine amide | (0.5 M) | Water | ArgMetNH ₂ | 38% |
| L-Phenylalanine amide | (0.2 M) | 30% EtOH | ArgPheNH ₂ | 6% |
| L-Phenylalanine amide | (0.6 M) | 30% DMF | ArgPheNH ₂ | 65% |

^{a)} 5 uM enzyme, 50 mM CaCl₂, 10 mM DTT, pH adjusted with TEA

Example 12

Porcine Pancrease Lipase ^{a)} catalyzed synthesis of L,L-dipeptide amides using L-amino acid ethyl esters (50 mM) as substrates and L-amino acid amides as nucleophiles in 0% EtOH at pH 8.5

| Substrate | Nucleophile | (conc.) | Product | Yield |
|-----------|------------------------------------|---------|-----------------------|-------------------|
| L-TrpOEt | L-MetNH ₂ | (1.5 M) | TrpMetNH ₂ | 71% |
| L-TrpOEt | L-MetNH ₂ | (1.0 M) | TrpMetNH ₂ | 54% |
| L-TyrOEt | L-SerNH ₂ | (1.5 M) | TrpSerNH ₂ | 69% |
| L-MetOEt | L-MetNH ₂ ^{b)} | (1.0 M) | MetMetNH ₂ | 31% ^{c)} |

^{a)} 500 uM

^{b)} Reaction in pure water

^{c)} At incomplete conversion

Example 13

Candida Cylindracea Lipase ^{a)} catalyzed synthesis of L,L-dipeptide amides using L-amino acid ethyl esters (50 mM) as substrates and L-amino acid amides as nucleophiles in 30% EtOH at pH 8.5

| Substrate | Nucleophile | (conc.) | Product | Yield |
|-----------|----------------------|---------|-----------------------|-------------------|
| L-TrpOEt | L-MetNH ₂ | (1.5 M) | TrpMetNH ₂ | 75% ^{b)} |
| L-TyrOEt | L-SerNH ₂ | (1.5 M) | TyrSerNH ₂ | 50% ^{b)} |

^{a)} 1000 uM, prolonged reaction time

^{b)} Versus hydrolysis at less than 50% conversion

Example 14

Preparative synthesis of L,L-Tryptophanymethionine amide, TrpMetNH₂

Procedure

L-Tryptophane ethyl ester hydrochloride (4.0 g, 15 mmol) and L-Methionine amide hydrochloride (55.7 g, 300 mmol) were dissolved in 195 ml H₂O and 90 ml ethanol, and pH was adjusted to 8.5 with sodium hydroxide. The reaction was initiated by addition of 0.8 g of crude Porcine Pancreatic Lipase and was kept at pH 8.5 for the duration of the reaction. The remainder of the substrate (4.0 g, 15 mmol) was added after 5 hours and the reaction continued overnight. It was then stopped by adjusting pH to 3 with HCl-solution.

The mixture was then diluted, ethanol was removed by evaporation under reduced pressure, and the mixture was filtered. The filtrate was purified by RP-preparative HPLC (Waters Prep LC/System 500A) using two columns (5.7 x 30 cm) packed with 60 um C-18 particles and 5 mM HCl/ethanol mixtures as eluent.

Collected fractions containing pure product were concentrated under reduced pressure and finally freeze dried.

This procedure gave 4.78 g of L,L-Tryptophanymethionine amide hydrochloride (12.9 mmol, 43%) as an amorphous powder.

Identification

The product was identified as the hydrochloride containing 10.3% (w.w) of chloride.

Amino acid analysis showed the absence of free amino acids and gave the following results after acid

5 hydrolysis:

Met (1.00)

Trp (1.00)

Specific optical rotation in 50% MeOH, $c = 0.2$ using the sodium D-line was found to be $+40.0^\circ$ at 20°C .

10

Purity

HPLC-purity: 92.4% (Novapak 4 μm C-18, 0.1 M ammonium phosphate, pH 3.0/acetonitrile, 220 nm)

Water content by Karl Fisher: 5.3% (w.w)

15 Quantization of the alpha-amino group by reaction with Trinitrobenzene sulphonic acid and UV-detection: 75.8% (w.w).

Peptide content by UV-quantization: 88.3% (w.w) (281 nm, Trp-absorbance in MeOH: 0.1 N KOH (1:1))

Example 15

20

Preparative synthesis of L,D-Tyrosylserin amide, TyrserNH₂

25

Procedure

L-Tyrosine ethyl ester hydrochloride (2.5 g, 10 mmol) and D-serine amide hydrochloride (17.5 g, 100 mmol) were dissolved in 200 ml of water, and pH was adjusted to 8.5 with sodium hydroxide. The reaction was initiated by addition of 50 mg of alpha-chymotrypsin and was kept at pH 8.5 for the duration of the reaction. After 30 minutes precipitation of free tyrosine began. The remainder of the substrate (5.0 g, 20 mmol) was added in two portions of 2.5 g during 1 hour. The reaction was stirred for two hours and was then stopped by adjusting pH to 3 with HCl-solution.

30 The formed tyrosine was filtered off, and the filtrate was purified by RP-preparative HPLC (Waters Prep LC-System 500A) using two columns (5.7 x 30 cm) packed with 20 μm C-18 particles and 50 μm acetic acid as an eluent.

Collected fractions containing pure product were concentrated under reduced pressure and finally freeze dried with addition of aqueous HCl.

40 This procedure gave 2.8 g of L,D-Tyrosylserine amide hydrochloride (9.2 mmol, 31%) as an amorphous powder.

Identification

45 The product was identified as the hydrochloride containing 16.0% (w/w) of chloride.

Amino acid analysis showed the absence of free amino acids, but following acid hydrolysis gave the following results:

Ser (1.10)

Tyr (0.90)

50 Specific optical rotation in 50% MeOH, $c = 0.3$ using the sodium D-line was found to be $+58.0^\circ$ at 20°C .

Purity

55 HPLC-purity: 97.4% (Novapak 4 μm C-18, 0.1 M ammonium phosphate, pH 3.0/acetonitrile, 220 nm)

Water content by Karl Fisher: 1.8% (w/w).

Quantization of the alpha-amino group by reaction with Trinitrobenzene sulphonic acid and UV-detection: 81% (w/w).

Example 16

5 Preparative synthesis of D,L-tyrosylLeucin amide, tyrLeuNH₂Procedure

10 D-tyrosine ethyl ester hydrochloride (3.5 g, 14 mmol) and L-Leucin amide hydrochloride (14 g, 84 mmol) were dissolved in 246 ml of 0.1 M KCl and pH was adjusted to 9.0 with sodium hydroxide. The reaction was initiated by addition of 0.7 g of alpha-chymotrypsin and stirred for two days at room temperature. pH was kept at 9.0 for the duration of the reaction. The reaction was stopped by adjusting pH to 3 using HCl-solution.

15 The formed tyrosine was filtered off, and the filtrate was purified by RP-preparative HPLC (Waters Prep LC-System 500A) using two columns (5.7 x 30 cm) packed with 20 uM C-18 particles and 5 mM HCl as an eluent.

Collected fractions containing pure product were concentrated by evaporation under reduced pressure and finally freeze dried.

20 This procedure gave 2.50 g of D,L-tyrosylLeucine amide hydrochloride (7.6 mmol, 54%) as an amorphous powder.

Identification

25 The product was identified as the hydrochloride containing 9.8% (w/w) of chloride.
Amino acid analysis showed the absence of free amino acids, but following acid hydrolysis gave the following results:

Tyr (0.98)

30 Leu (1.03)

Specific optical rotation in water, c=0.1 using the sodium D-line was found to be -129.4° at 20°C.

Purity

35 HPLC-purity: 92.9% (Novapak 4 um C-18, 0.1 M ammonium phosphate, pH 3.0/acetonitrile, 220 nm)
Water content by Karl Fisher: 6.8% (w/w).

Quantization of the alpha-amino group by reaction with Trinitrobenzen sulfonic acid and UV-detection: 74.9%

40 UV-quantization: 74.9% (Tyrosine phenolate absorbance at 293 nm, in 0.1 M KOH)

Example 17

45 Preparative synthesis of L,L-ArginylMethionine amide, ArgMetNH₂Procedure

50 L-Arginine ethyl ester dihydrochloride (4.1 g, 15 mmol) and L-Methionine amide hydrochloride (55.4 g, 300 mmol) were dissolved in 300 ml of water, and pH was adjusted to 8.5 with sodium hydroxide. The reaction was initiated by addition of 50 mg of trypsin. pH was kept at 8.5 for the duration of the reaction. The remainder of the substrate (8.2 g, 30 mmol) was added during one hour. The reaction was then
55 stopped by adjusting pH to 3 using HCl-solution.

The reaction mixture was then diluted and purified by successive cation exchange on a DOWEX A650 Wx4 and a CM-Sephacrose 6B column using ammonium acetate and NaCl/HCl salt gradients, respectively, and was finally desalted.

Collected fractions containing pure product were concentrated under reduced pressure and finally freeze dried.

This procedure gave 10.7 g of L,L-ArginylMethionine amide dihydrochloride (28.3 mmol, 63%) as a white amorphous powder.

Identification

Less than 0.2% (w/w) of acetate and 22.9% (w/w) of chloride were measured, so the product was present as a dihydrochloride.

Amino acid analysis showed the absence of free amino acids, but following acid hydrolysis gave the following results:

Tyr (1.00)

Leu (0.80)

Specific optical rotation in 50% MeOH, $c = 0.2$ using the sodium D-line was found to be $+19.5^\circ$ at 20°C .

Purity

HPLC-purity: 95.1% (Novapak C-18, 0.1 M ammonium phosphate containing alkylsulfonate, pH 4.5/acetonitrile, 220 nm)

Water content by Karl Fisher: 9.1% (w/w).

Peptide content by amino acid analysis: 72% (w/w) based on Arginine.

| Example 18 | | | | |
|--|-------------|----|-----------------|---------------------|
| Synthesis of L,L-Methionyl-Methionine amide catalyzed by Eupergit C immobilized Porcine Pancreatic Lipase ^{a)} using L-Methionine ethyl ester as substrate and L-Methionine amide as nucleophile at pH 8.5 in water and water/organic homogeneous mixtures. | | | | |
| Conc. of substrate | Nucleophile | % | Organic solvent | Yield ^{b)} |
| 50 mM | 1.0 M | 0 | - | 68% |
| 100 mM | 0.5 M | 30 | Isopropanol | 26% |

^{a)} Porcine Pancreatic Lipase was immobilized on Eupergit C using the procedure recommended by the manufacturer to a final concentration corresponding to approx. 400 μM on the gel by activity assay.

^{b)} The reaction mixtures with volumes of 1-3 reaction bed volumes were recirculated over the column packed with enzyme gel until full conversion of the substrate as determined by HPLC (0.5-2 days) while pH was being kept constant by pH-stat control.

Example 19

Porcine Pancreatic Lipase ^{a)} and alpha-chymotrypsin ^{a)} catalyzed synthesis of L,L-Tryptophanyl-Alanine-tert-butylester in pure organic solvents with L-Tryptophane ethyl ester (50 mM) as substrate and L-Alanine-tert-butylester (0.3 M) as nucleophile, both in saltfree form.

| Enzyme ^{a)} | Solvent | Conversion ^{c)} | Yield ^{c)} |
|----------------------|--|--------------------------|---------------------|
| PPL | CH ₂ Cl ₂ /n-Hexane (1:3) | 30% | 13% |
| CT | CH ₂ Cl ₂ /n-Hexane (1:3) | 56% | 17% |
| PPL | CH ₂ Cl ₂ | 7% | 10% |
| CT | CH ₂ Cl ₂ | 50% | 3% |
| PPL ^{b)} | CH ₂ Cl ₂ /Isooctane (1:1) | 82% ^{b)} | 26% |

^{a)} 500-2000 μ M freeze-dried enzyme containing moisture was added directly to the mixture, which was stirred for 24 hours.

^{b)} In this case, 500 μ M enzyme and 3 days of reaction time were applied.

^{c)} Conversion of starting material and yield versus hydrolysis was determined by HPLC of samples quenched with DMF, evaporated and redissolved in DMF/acid water. Controls showed neither conversion nor yield.

Example 20

Alpha-chymotrypsin ^{a)} catalyzed synthesis of sidechain protected L,L-dipeptide amides with L-Tyrosine or L-Phenylalanine ethyl esters (50 mM) as substrates and L-S-Acetyl-L-methylcysteine amide (0.6 M) as nucleophiles in water at pH 8.5.

| Substrate | Product | Yield |
|-----------------------------|------------------------------|-------|
| L-Tyrosine ethyl ester | TyrCys(-SACm)NH ₂ | 62% |
| L-Phenylalanine ethyl ester | PheCys(-SACm)NH ₂ | 78% |

^{a)} 5 μ M

Example 21

Alpha-chymotrypsin ^{a)} catalyzed synthesis of L,L-Dipeptide alcohols using L-Tyrosine and L-Phenylalanine ethyl esters (50 mM) as substrates and L-Amino acid alcohols (0. M) as nucleophiles in water at pH 8.5.

| Substrate | Nucleophile | Product | Yield |
|-----------|-------------------------|--------------------------|-------|
| L-TyrOEt | L-MetCH ₂ OH | TyrMetCH ₂ OH | 42% |
| L-TyrOEt | L-LeuCH ₂ OH | TyrLeuCH ₂ OH | 46% |
| L-PheOEt | L-MetCH ₂ OH | PheMetCH ₂ OH | 60% |

^{a)} 10 μ M

Example 22

Synthesis of L,L-dipeptides catalyzed by thiolendoproteases ^{a)} using amino acid ethyl esters (50 mM) as substrate and free L-Amino acids as nucleophiles in water at pH 8.5.

| Enzyme | Substrate | Nucleophile | (conc.) | Product | Yield ^{b)c)} |
|--------|-----------|-------------|---------|----------|-----------------------|
| Ficin | ArgOEt | ArgOH | (1.0 M) | ArgArgOH | 6% |
| Papain | ArgOEt | ArgOH | (1.0 M) | ArgArgOH | 7% |
| Ficin | LysOEt | AlaOH | (1.5 M) | LysAlaOH | 5% |
| Papain | LysOEt | AlaOH | (1.5 M) | LysAlaOH | 6% |

^{a)} 100 μ M, 2 mM EDTA, 0.1 M KCl, 5 mM DTT or 10 mM cysteine.

^{b)} Determined vs. hydrolysis via a standard at less than 50% conversion.

^{c)} Controls without enzyme showed no detectable aminolysis under the conditions reported.

Example 23

Trypsin ^{a)} catalyzed synthesis of L,L-dipeptide amides with Arginine-paranitroanilide (10 mM) as substrate and L-Amino acid amides (0.3 M) as nucleophiles in 40% DMF at H 8.5.

| Nucleophile | Product | Yield ^{b)} |
|------------------|-----------------------|---------------------|
| Methionine amide | ArgMetNH ₂ | 45% |
| Leucine amide | ArgLeuNH ₂ | 31% |
| Tyrosine amide | ArgTyrNH ₂ | 14% |

^{a)} 5 μ M

^{b)} Determined vs. hydrolysis via a standard at less than 80% conversion.

Example 24

Papain ^{a)}^{b)}
catalyzed
synthesis of
L,L-Lysyl-Alanine
amide using
L-Lysine
ethyl ester
(50 mM) as
substrate
and
L-Alanine
amide (0.8
M) as
nucleophile
in water at
various
pH-values.

| pH | Yield ^{c)} |
|-----|---------------------|
| 6.5 | 21% |
| 4.5 | 47% |

^{a)} 50 μ M, 2 mM EDTA, 10 mM Cysteine.

^{b)} At prolonged reaction time; less than 50% conversion.

^{c)} Determined vs. hydrolysis using a standard and corrected for incomplete conversion.

Example 25

Alpha-chymotrypsin ^{a)} and clostripain ^{b)} catalyzed synthesis of L,L-dipeptide esters using L-Amino acid ethyl esters (50 mM) as substrates and L-Amino acid ethyl or tert-butyl-esters as nucleophiles in water at pH 7.5 and 8.5.

| Enzyme | Substrate | Nucleophile | (conc.) | Product | Yield |
|--------|-----------|-------------|---------|------------|-------------------|
| CT | TrpOEt | AlaOtBu | (0.8 M) | TrpAlaOtBu | 12% |
| CT | TrpOEt | ValOEt | (0.8 M) | TrpValOEt | 18% |
| CL | ArgOEt | MetOEt | (1.0 M) | ArgMetOEt | 33% ^{c)} |

^{a)} 5 μ M, 0.1 M KCl, pH 7.5.

^{b)} 10 μ M, 50 mM CaCl₂, 10 mM DTT, pH 8.5 adjusted with TEA.

^{c)} Determined at less than 50% conversion.

Example 26

Porcine Pancreatic Lipase ^{a)} and Lipolase (Novo)(LIP) ^{a)} and Rhizopus Arrhizus Lipase (RA) ^{b)} catalyzed synthesis of L,L-Methionyl-Methionine amide using different L-Methionine esters as substrates and L-Methionine amide as nucleophile in various aqueous/organic solvent homogeneous mixtures at pH 8.5 and 1.0 M nucleophile, unless otherwise indicated.

| Enzyme | Ester substrate | (conc.) | % | Organic solvent | Yield ^{e)} |
|-------------------|-----------------|----------|----|------------------|---------------------|
| PPL ^{c)} | Ethyl | (50 mM) | 0 | - | 55% |
| PPL ^{d)} | Ethyl | (50 mM) | 0 | - | 36% |
| PPL | Ethyl | (150 mM) | 0 | - | 42% |
| PPL | Ethyl | (200 mM) | 15 | Dimethoxy Ethane | 20% |
| PPL | Ethyl | (150 mM) | 30 | Isopropanol | 36% |
| PPL | Ethyl | (50 mM) | 90 | Ethylene Glycol | 42% |
| PPL | n-Propyl | (50 mM) | 0 | - | 55% |
| PPL | n-Hexyl | (50 mM) | 0 | - | 39% |
| LIP | Ethyl | (50 mM) | 0 | - | 47% |
| LIP | Ethyl | (50 mM) | 15 | Dimethoxy Ethane | 35% |
| LIP | n-Hexyl | (50 mM) | 0 | - | 32% |
| RA | Ethyl | (50 mM) | 30 | Ethanol | 23% |

^{a)} 1000 μ M - 2000 μ M

^{b)} 50 μ M

^{c)} pH 9.0

^{d)} At 0.25 M nucleophile

^{e)} Various minor amounts of deamidation of product was noted under these conditions and is not included in the yields reported.

Example 27

Trichoderma Viridae Cellulase catalyzed synthesis of L,L-TyrGlyNH₂ using L-Tyrosine ethyl ester as a substrate and Glycine amide as nucleophile in water.

| Substrate | (conc.) | Nucleophile | (conc.) | Product | Yield ^{c)} |
|----------------------|---------|--------------------|---------|-----------------------|---------------------|
| TyrOEt ^{a)} | (20 mM) | GlyNH ₂ | (0.6 M) | TyrGlyNH ₂ | 23% |
| TyrOEt ^{b)} | (10 mM) | GlyNH ₂ | (0.8 M) | TyrGlyNH ₂ | 34% |

^{a)} 1000 μ M crude enzyme, pH 8.5

^{b)} 500 μ M crude enzyme, pH 8.0

^{c)} Determined vs. hydrolysis at less than 30% conversion and corrected for hydrolysis and aminolysis found in controls at similar conditions.


Claims

1. A process for producing dipeptides or structurally related compounds having the general formula H-A-B-Y

wherein A represents an optionally side-chain protected L- or D- α -amino acid residue or ω -amino acid residue and B represents an optionally side-chain protected L- or D- α -aminocarboxylic acid residue which may be the same as or different from A, an L- or D-aminophosphonic acid residue or L- or D-aminosulfonic acid residue or the corresponding ω -amino acids or salts and hydrates thereof, and Y is OH, H, alkyl, aryl, aralkyl or a C-terminal blocking group, or BY represents an amino alcohol residue

B'-CHY'-OH

wherein B' is a decarboxy derivative of the aminocarboxylic acids as defined with relation to B, and Y' is H, alkyl, aryl or aralkyl, characterized by reacting a substrate component, which is an amino acid derivative having the formula



wherein A is as defined above, R¹ represents alkyl, aryl or aralkyl optionally substituted with inert substituents or an α -des-amino fragment of an amino acid, and R² and R³ are the same or different and each represents hydrogen, alkyl, aryl or aralkyl optionally substituted with inert substituents,


with a nucleophile component selected from

(a) amino acids having the formula

H-B-OH

wherein B is an aminocarboxylic acid residue as defined above,

(b) amino acid amides having the formula



wherein B is an aminocarboxylic acid residue as defined above, and R² and R³ have the above meaning, except that when R² represents hydrogen, R³ may also represent hydroxy or amino,

(c) amino acid esters having the formula

H-B-OR⁴

wherein B is an aminocarboxylic acid residue as defined above, and R⁴ represents alkyl, aryl or aralkyl,

(d) optionally acid group protected straight chain or branched aminophosphonic acids or aminosulfonic acids having the formula

NH₂C_xH₂PO₃R⁵R⁶ or NH₂C_xH₂SO₃R⁷

wherein R⁵, R⁶ and R⁷ independently represent hydrogen, alkyl, aryl or aralkyl, x is 1-6 and z is 2-12,

(e) amino acid aldehydes or ketones or derivatives thereof having the formula

H-B'-CY²-R⁸

wherein B' is as defined above, Y² is O or a functional derivative thereof, preferably a ketal, and R⁸ is H, alkyl, aryl or aralkyl, and

(f) amino alcohols having the formula

H-B'-CHY'-OH

wherein B' and Y' have the above meaning

in the presence of an amidase or esterase enzyme different from serine or thiol carboxypeptidases in solution or dispersion, and then, if desired, cleaving a present side-chain protecting group or protective group Y and/or, if desired, converting the resulting dipeptide derivative to a salt or hydrate.

2. A process according to claim 1, characterized by using an enzyme with esterase or amidase activity selected from serine or thiolendoproteases, lipases, esterases and glycosidases.

3. A process according to claim 1 or 2, **characterized** in that the enzyme used has been chemically modified or is a biosynthetic mutant of a natural form.

4. A process according to any of the preceding claims, **characterized** by using an immobilized enzyme.

5 5. A process according to any of the preceding claims, **characterized** by using an aqueous reaction solution or dispersion containing 0-90%, preferably 0-60%, of a polar water miscible organic solvent and having a pH value of 3-11, preferably 5-10.5, more preferably 6-10, in particular 7-9.5.

6. A process according to claim 5, **characterized** in that the organic solvent is selected from alkanols, dimethyl sulfoxide, dimethyl formamide, dimethoxy ethane and ethylene glycol.

10 7. A process according to any of claims 1 to 4, **characterized** by using an organic reaction solution or dispersion containing 0-10% of water.

8. A process according to claim 7, **characterized** by using an unpolar organic solvent, preferably selected from dialkyl ethers, ethyl acetate, ethyl propionate, octanes, heptanes, hexanes, petroleum ether and methylene chloride.

15 9. A process according to claim 7, **characterized** by using a liquid substrate or nucleophile component which may also serve as the organic solvent.

10. A process according to any of the preceding claims, **characterized** by using as substrate component a D- or L-amino acid ester selected from benzyl esters or straight or branched C₁-C₈ alkyl esters optionally substituted with inert substituents.

20 11. A process according to any of the preceding claims, **characterized** by using as nucleophile component an amino acid amide having the formulae
H-B-NHR³

wherein R³ is hydrogen or C₁-C₃ alkyl and B is an L- or D-amino carboxylic acid residue.

25 12. A process according to any of claims 1 to 10, **characterized** by using as nucleophile component an ester having the formula
H-B-OR⁴

wherein B is a L- or D-aminocarboxylic acid residue and R⁴ is C₁-C₃ alkyl.

30 13. A process according to any of the preceding claims, **characterized** in that the resulting dipeptide includes one or more C-terminal protective groups Y, and that the group or groups are cleaved enzymatically, either by means of the same enzyme as was used in the preceding reaction or by means of an enzyme having a different ester or amide specifically.

14. A process according to any of the preceding claims, **characterized** in that the resulting dipeptide includes one or more side-chain protective groups and that the group or groups are cleaved enzymatically, preferably by means of an esterase or lipase or proteolytical enzyme.

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

EP 89 30 8232.1

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|---|---|--|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int. Cl. 4) |
| Y | EP-A2-0 269 390 (GENENCOR INC.) *See page 2 lines 33-60* | 1-14 | C 12 P 21/02 C 07 K 1/00 |
| | --- | | |
| Y,D | EP-A2-0 220 923 (W.R. GRACE & CO) *See claim 1* | 1-14 | |
| | --- | | |
| Y,D | EP-A1-0 074 095 (TOYO SODA MANUFACTURING CO., LTD.) | 1-14 | |
| | --- | | |
| Y | EP-A2-0 154 472 (AJINOMOTO CO., INC.) | 1-14 | |
| | --- | | |
| Y,D | EP-A1-0 017 485 (DE FORENEDE BRYGGERIER A/S) *See page 5, lines 1-5, pages 6-7 and the claims* | 1-14 | TECHNICAL FIELDS SEARCHED (Int. Cl. 4) C 07 K C 12 P |
| | --- | | |
| P,D | EP-A1-0 278 787 (CARLSBERG BIOTECHNOLOGY LTD. A/S) | 1-14 | |
| The present search report has been drawn up for all claims | | | |
| Place of search STOCKHOLM | | Date of completion of the search 09-10-1989 | Examiner SIÖSTEEN Y. |
| <p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p> | | | |